

Middlesex University Research Repository

An open access repository of

Middlesex University research

<http://eprints.mdx.ac.uk>

Paice, Katherine, Dimitriou, Lygeri ORCID logoORCID: <https://orcid.org/0000-0002-5093-558X>,
Wilson, Laura J. ORCID logoORCID: <https://orcid.org/0000-0002-3744-6132>, Hills, Frank
ORCID logoORCID: <https://orcid.org/0000-0001-8235-7545>, Sinclair, Scott, Gondek, Marcela
B., Kwan, Xue Bin and Wood, Alyssa (2017) Influence of cold water immersion and cherry juice
on recovery following a 42.2km trail run. BASES Conference 2017 – Programme and Abstracts
(Journal of Sports Sciences, Vol. 35, Iss. sup1, 2017). In: BASES Conference 2017, 28-29 Nov
2017, Nottingham, UK. . ISSN 0264-0414 [Conference or Workshop Item]
(doi:10.1080/02640414.2017.1378421)

Final accepted version (with author's formatting)

This version is available at: <https://eprints.mdx.ac.uk/23226/>

Copyright:

Middlesex University Research Repository makes the University's research available electronically.

Copyright and moral rights to this work are retained by the author and/or other copyright owners unless otherwise stated. The work is supplied on the understanding that any use for commercial gain is strictly forbidden. A copy may be downloaded for personal, non-commercial, research or study without prior permission and without charge.

Works, including theses and research projects, may not be reproduced in any format or medium, or extensive quotations taken from them, or their content changed in any way, without first obtaining permission in writing from the copyright holder(s). They may not be sold or exploited commercially in any format or medium without the prior written permission of the copyright holder(s).

Full bibliographic details must be given when referring to, or quoting from full items including the author's name, the title of the work, publication details where relevant (place, publisher, date), pagination, and for theses or dissertations the awarding institution, the degree type awarded, and the date of the award.

If you believe that any material held in the repository infringes copyright law, please contact the Repository Team at Middlesex University via the following email address:

eprints@mdx.ac.uk

The item will be removed from the repository while any claim is being investigated.

See also repository copyright: re-use policy: <http://eprints.mdx.ac.uk/policies.html#copy>

Influence of cold water immersion and cherry juice on recovery following a 42.2km trail run.

Paice, K.¹, Dimitriou, L.¹, Wilson, L.¹, Hills, FA.¹, Sinclair, S.¹, Gondek, MB.¹, Kwan Xue Bin, K.¹, Wood, A.¹, and Cockburn, E.¹

¹Middlesex University (Middlesex, UK)

*Corresponding author: Kat Paice (kat.paice@outlook.com); @kat_paice

Both cherry juice (CJ) (Howatson et al, 2009, *Scandinavian Journal of Medicine and Science in Sport*, 20, 843-852) and cold water immersion (CWI) (White and Wells, 2013, *Extreme Physiology and Medicine*, 2) are effective for recovery following strenuous exercise. Therefore, this study examined whether CWI in conjunction with CJ is more effective for recovery than either alone. With ethical approval 38 endurance-trained males (age 40 ± 7 years; height 1.8 ± 7.7 m; mass 76.2 ± 10.0 kg) were randomly assigned to one of four interventions (placebo (PL); CJ; CWI; CWI and CJ (MIX)), before completing a 42.2km run. Markers of muscle damage (muscle soreness (DOMS); maximum voluntary isometric contraction (MVIC); creatine kinase (CK)), inflammation (interleukin-6 (IL-6); C-reactive protein (CRP)) and oxidative stress (protein carbonyls (PC); superoxide dismutase (SOD); Thiobarbituric acid reactive substances (TBARS)) were measured at baseline (B), and immediately (P) (IL-6, PC and SOD only), 24h and 48h post run. Probabilistic magnitude based inferences analyses (difference; $\pm 90\%CI$; difference; $x/\div 90\%CI$ for blood variables which are reported as factors), for changes over time, revealed MIX was *likely harmful* versus PL (-11 ; $\pm 7\%$), CWI (-9 ; $\pm 7\%$) and CJ (-8 ; $\pm 8\%$) for changes in MVIC B-24h. MIX was *likely*, *possibly* and *likely harmful* versus PL (1.4 ; $x/\div 1.5$), CWI (1.2 ; $x/\div 1.7$) and CJ (1.7 ; $x/\div 1.6$) for CK B-48h. B-24h MIX was *possibly*, *most likely* and *very likely harmful* for changes in CRP versus PL (1.8 ; $x/\div 2.7$), CWI (8.2 ; $x/\div 2.3$) and CJ (5.4 ; $x/\div 2.6$). B-post MIX was *likely harmful* (2.8 ; $x/\div 2.4$) and *possibly harmful* (2.0 ; $x/\div 2.4$) in comparison to PL and CWI for changes in IL-6. B-48h MIX was *likely*, *likely* and *very likely harmful* in comparison to PL (1.4 ; $x/\div 1.5$), CWI (1.8 ; $x/\div 1.6$) and CJ (2.1 ; $x/\div 1.5$) for changes in SOD. B-24h MIX was *likely* (0.5 ; $x/\div 1.8$) and *possibly* (0.7 ; $x/\div 1.8$) *beneficial* for TBARS in comparison to PL and CJ. Independently both treatments have been found to improve recovery; together they may limit blood supply to damaged tissues and reduce some oxidative stress markers which may be necessary for an accelerated recovery. However, further investigation is needed to clarify this and any other potential mechanisms. In conclusion, combined use of CWI and CJ is detrimental to recovery, based on measures of muscle function, inflammation and oxidative stress, following a 42.2km run.